



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07K 3/02, 7/10, 15/24 C12P 21/06, A23J 1/22, 3/10	A1	(11) International Publication Number: WO 92/18526 (43) International Publication Date: 29 October 1992 (29.10.92)
(21) International Application Number: PCT/AU92/00175 (22) International Filing Date: 16 April 1992 (16.04.92) (30) Priority data: PK 5706 19 April 1991 (19.04.91) AU (71) Applicants (for all designated States except US): THE UNIVERSITY OF MELBOURNE [AU/AU]; Grattan Street, Parkville, VIC 3052 (AU). THE VICTORIAN DAIRY INDUSTRY AUTHORITY [AU/AU]; 651-653 Victoria Street, Abbotsford, VIC 3067 (AU). (72) Inventor; and (75) Inventor/Applicant (for US only) : REYNOLDS, Eric, Charles [AU/AU]; School of Dental Science, The University of Melbourne, Royal Dental Hospital of Melbourne, 711 Elizabeth Street, Melbourne, VIC 3000 (AU).		(74) Agent: CARTER SMITH & BEADLE; Qantas House, 2 Railway Parade, Camberwell, VIC 3124 (AU). (81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), MC (European patent), NL (European patent), SE (European patent), US. Published <i>With international search report.</i>
(54) Title: PRODUCTION OF PHOSPHOPEPTIDES FROM CASEIN (57) Abstract <p>A method for the preparation of selected phosphopeptides having anticariogenic and other activities, comprising the steps of completely digesting a soluble monovalent cation salt of casein in solution with a proteolytic enzyme, adding a mineral acid to the solution to adjust the pH to about 4.7, removing any precipitate produced, adding CaCl₂ to a level of about 1.0 % w/v to cause aggregation of at least the selected phosphopeptides in said digested solution, separating the aggregated phosphopeptides from the solution through a filter having a molecular weight exclusion limit lying substantially within the range 10,000 to 20,000 while passing the bulk of the remaining phosphopeptides and solution, diafiltering the separated phosphopeptides with water through a filter and concentrating and drying the retentate.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
DE	Germany	MC	Monaco	TC	Togo
DK	Denmark			US	United States of America

- 1 -

TITLE: PRODUCTION OF PHOSPHOPEPTIDES FROM CASEIN**Field of the Invention:**

This invention relates to the production of phosphopeptides having anticariogenic and other properties from casein.

Background of the Invention:

In our Australian Patent No. 593365, we have described that four of the many phosphopeptides released by tryptic digestion of casein have anticariogenic (tooth-decay-inhibiting) activity. These peptides all contain the active sequence - Ser(P)-Ser(P)-Ser(P)-Glu-Glu- and correspond to α_{11} (59-79) SEQ.ID No 2 (T_1), B(2-25) SEQ.ID No 3 (T_2), α_{12} (46-70) SEQ.ID No 4 (T_4) and α_{12} (2-21) SEQ.ID No 7 (T_3). The methods described for the production of the anticariogenic phosphopeptides are selective precipitation and ion exchange chromatography. While these methods produce very pure preparations of these peptides, they have not received general acceptance in the dairy industry due to their cost and the level of technical skill required.

Recently membrane ultrafiltration has found broad acceptance in the dairy industry for milk treatment. In U.S. Patents 4,358,465, 4,361,587 and 4,495,176, Brule et al describe an ultrafiltration method for the

- 2 -

production of casein phosphopeptides as dietetic aliments. This procedure proves unsuitable for the production of anticariogenic phosphopeptides due to the predominance of non-anticariogenic phosphopeptides in the preparations.

Summary of the Invention and Object:

It is an object of the present invention to provide a method of preparing selected phosphopeptides from casein using ultrafiltration.

The invention provides a method for the preparation of selected phosphopeptides comprising the steps of completely digesting a soluble monovalent cation salt of casein in solution, introducing a di or trivalent metal ion to cause aggregation of at least the selected phosphopeptides in said digested solution, and filtering the solution through a filter having a molecular weight exclusion limit selected to retain at least said aggregated phosphopeptides while passing the bulk of the remaining phosphopeptides.

In the method described by Brule et al, the object is to obtain a broad range of phosphopeptides from casein for use as a dietetic aliment. Therefore Brule et al do not teach that the casein compound must be completely hydrolysed or that a di/trivalent metal ion must be added to aggregate the desired phosphopeptides to a size which enables those

- 3 -

phosphopeptides to be filtered from the solution while allowing the remaining phosphopeptides and solution to pass. This represents a significant advance in the art since it enables the use of an industry accepted method of extraction which results in a preparation which is rich (> 90% w/w) in the desired phosphopeptides.

In a preferred form of the invention, the selected phosphopeptides are the anticariogenic phosphopeptides referred to above, and the molecular exclusion limit adopted during the filtering step of the above method preferably substantially falls within the range 10,000 to 20,000.

The soluble monovalent cation salt of casein, such as sodium caseinate or potassium caseinate, may be present in the solution in a concentration substantially falling within the range 0.1 to 50% w/w, which is preferably digested using a proteolytic enzyme, such as pancreatin, trypsin, papain or chymotrypsin, or a mixture of proteolytic enzymes such as trypsin and chymotrypsin or by chemical means, such as cyanogen bromide. The enzyme(s) to casein ratio can range from about 1:1000 to 1:10 (w:w) but this would be selected to allow complete digestion of the casein as defined above. The pH of the hydrolysis should preferably be controlled at optimum for the enzymes to

- 4 -

allow complete casein digestion. The temperature also should be optimised for complete digestion but temperature induced degradation (deamidation, dephosphorylation and peptidolysis) should be minimised. The optimal temperature is between about 20°C and 60°C.

In a preferred form of the invention, after digestion HCl is added at room temperature to about pH 4.7 and any precipitate (this should be minimal) removed. CaCl₂ is then added to the supernatant to a level of about 1.0% w/v. Phosphopeptides in the presence of 1.0% w/v calcium (II) aggregate. The anticariogenic phosphopeptides (ie. containing the sequence -Ser(P)-Ser(P)-Ser(P)-Glu-Glu-) form hexamers which are separated from the smaller non-anticariogenic phosphopeptide aggregates by extensive diafiltration through a 10,000 molecular weight exclusion limit filter with a CaCl₂ solution preferably 1.0% w/v.. The preferred molecular weight exclusion limit of the membrane filter should not be less than 10,000 or greater than about 20,000. The addition of a CaCl₂ solution, or some other suitable di/trivalent metal ion, such as zinc (II) or ferric (III), is essential for diafiltration in order to maintain the integrity of the anticariogenic phosphopeptide aggregates thus allowing separation of the

- 5 -

anticariogenic from the non-anticariogenic phosphopeptides.

After several volumes of 1.0% CaCl_2 w/v have passed through the membrane filter to achieve greater than 90% purity of the anticariogenic phosphopeptides the ultraretentate containing the anticariogenic phosphopeptides can be diafiltered with water through a 1,000 molecular weight exclusion limit filter to remove calcium if desired. The retentate is then concentrated and spray dried.

The calcium, zinc and ferric salts of the anticariogenic phosphopeptide preparation (ACPP) can be converted to a sodium salt by acidifying a 10% w/v solution of the calcium ACPP to a low pH, circa pH 2.0, with HCl. After extensive diafiltration through a 1,000 molecular weight exclusion limit filter the retentate is neutralised to pH 7.0 with NaOH and then diafiltered with water through the same filter to remove excess sodium chloride.

The calcium ACPP can be converted to calcium phosphate ACPP by addition of CaCl_2 and Na_2HPO_4 where the Ca/P final ratio is 1.67. The peptide α_{11} (59-79) can bind 21 Ca and 13 PO_4 . The filtrate of the above process is suited for the purification of other bioactive casein peptides by size and charge-based separation technologies and can be used as

- 6 -

microbiological growth media, as dietary supplements after debittering or as a nitrogen fertilizer.

A presently preferred embodiment of the invention will now be described with reference to the following example.

EXAMPLE

Sodium caseinate was prepared by acidifying milk with 0.1 M HCl to pH 4.7 and neutralising the precipitate with NaOH to pH 7.0. A 10% w/v solution of sodium caseinate was prepared and adjusted to pH 8.0. Trypsin (Novo) was added to 0.2% w/v and the hydrolysis allowed to proceed to completion at 50°C with adjustment to pH 8.0 by constant addition of NaOH. The pH of the solution was then adjusted to pH 4.7 with 5 M HCl and the precipitate removed at room temperature by centrifugation. The supernatant was microfiltered through an 8 micron filter, and then adjusted to pH 7.0 with NaOH and CaCl₂ added to a level of 1.0% w/v. This solution was then diafiltered through an Amicon YM10 (10,000 molecular weight exclusion limit) with 3 to 5 volumes of 1.0% w/v CaCl₂. The retentate was then washed with 1 volume of distilled/deionised water through an Amicon YM1 filter (1,000 molecular weight exclusion limit). The individual peptides of this preparation were separated by ion exchange FPLC and reverse phase HPLC as described in the aforementioned patent and identified

- 7 -

by amino acid composition and sequence analyses after conversion of the Ser(P) residues to S-ethyl cysteine. An analysis of the preparation is shown in Table 1.

Table 1. Composition of an Anticariogenic Phosphopeptide Preparation

<u>Peptide*</u>	<u>w/w</u>
$\alpha_2(1-21)$ (SEQ.ID No 8)	0.8
B(1-25) (SEQ.ID No 1)	22.3
$\alpha_2(2-21)$ (SEQ.ID No 7) (T_1)	5.7
B(2-25) (SEQ.ID No 3) (T_2)	17.9
$\alpha_1(59-79)$ (SEQ.ID No 2) (T_1)	21.4
Desamido ⁷⁴ $\alpha_1(59-79)$ (SEQ.ID No 5)	6.3
$\alpha_2(46-70)$ (SEQ.ID No 4) (T_4)	6.8
Desamido ^{74,78} $\alpha_1(59-79)$ (SEQ.ID No 6)	6.4
$\alpha_1(43-79)$ (SEQ.ID No 9)	3.3
NAP*	9.1

* NAP = non-anticariogenic peptides

This preparation contains the four anticariogenic phosphopeptides described in the aforementioned patent, [B(2-25), T_1 , $\alpha_1(59-79)$, T_1 , $\alpha_2(2-21)$, T_2 and $\alpha_2(46-70)$, T_4], those related peptides incompletely hydrolysed by trypsin [$\alpha_2(1-21)$, B(1-25) and $\alpha_1(43-79)$] and also minor levels of the two deamidated forms of $\alpha_1(59-79)$, desamido⁷⁴ and desamido^{74,78} which result from temperature induced deamidation, this occurs in


- 8 -

an even greater extent in commercial production of sodium caseinate due to higher temperatures and extremes of pH, although the presence of the deamidated forms has no effect on anticariogenic activity. The anticariogenic phosphopeptides were 90.9% w/w of the peptides produced.

If pure α_{11} -casein is used in place of casein then α_{11} (59-79) will be obtained by this process with minor amounts of the deamidated forms of this peptide depending on hydrolysis conditions. If pure β -casein is used then only β (1-25) and β (2-25) will be obtained using this process.

When crude enzymes are used (such as pancreatin), slight truncation (both N- and C- terminally) of the nine listed phosphopeptides can occur. As long as this truncation is only slight, there is no loss of activity. In fact, pancreatin produces a preparation with slightly greater specific activity on a weight basis when compared with purified trypsin.

The sequences of the nine peptides, which include the peptides T₁ to T₉ of the aforementioned patent, and the other peptides referred to above are detailed in the following sequence listing



- 9 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: REYNOLDS, ERIC CHARLES
- (ii) TITLE OF INVENTION: PRODUCTION OF PHOSPHOPEPTIDES FROM CASEIN
- (iii) NUMBER OF SEQUENCES: 9
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE:
 - (B) STREET:
 - (C) CITY:
 - (D) STATE:
 - (E) COUNTRY:
 - (F) ZIP: ~
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: FLOPPY DISK
 - (B) COMPUTER: IBM PC COMPATIBLE
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: WORD PERFECT
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A)
 - (B)
 - (C)
- (ix) TELECOMMUNICATION INFORMATION:
 - (A)
 - (B)
 - (C)

(2) INFORMATION FOR SEQ.ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25
 - (B) TYPE: Amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Protein
- (ix) FEATURE:
 - (A) NAME/KEY: Phosphoserine
 - (B) LOCATION: 15
 - (D) OTHER INFORMATION: Post-translationally phosphorylated serine
- (ix) FEATURE:
 - (A) NAME/KEY: Phosphoserine
 - (B) LOCATION: 17
 - (D) OTHER INFORMATION: Post-translationally phosphorylated serine
- (ix) FEATURE:
 - (A) NAME/KEY: Phosphoserine
 - (B) LOCATION: 18
 - (D) OTHER INFORMATION:

-10-

- Post-translationally phosphorylated serine
- (ix) FEATURE:
 (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 19
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine

(xi) SEQUENCE DESCRIPTION: SEQ.ID NO:1:

Arg Glu Leu Glu Glu Leu Asn Val Pro Gly Glu Ile Val Glu Ser Leu
 1 5 10 15

Ser Ser Ser Glu Glu Ser Ile Thr Arg
 20 25

(2) INFORMATION FOR SEQ.ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21
 (B) TYPE: Amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Protein
- (ix) FEATURE:
 (A) NAME/KEY: Pyroglutamate
 (B) LOCATION: 1
 (D) OTHER INFORMATION:
 A certain amount will exist in this form
- (ix) FEATURE:
 (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 6
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine
- (ix) FEATURE:
 (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 8
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine
- (ix) FEATURE:
 (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 9
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine
- (ix) FEATURE:
 (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 10
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine
- (ix) FEATURE:
 (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 17
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine

(xi) SEQUENCE DESCRIPTION: SEQ.ID NO:2:

Gln Met Glu Ala Glu Ser Ile Ser Ser Ser Glu Glu Ile Val Pro Asn
 1 5 10 15

Ser Val Glu Gln Lys
 20

(2) INFORMATION FOR SEQ.ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24
 (B) TYPE: Amino acid
 (C) STRANDEDNESS: single

- // -

- (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: Protein
 (ix) FEATURE:
 (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 14
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine
 (ix) FEATURE:
 (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 16
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine
 (ix) FEATURE:
 (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 17
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine
 (ix) FEATURE:
 (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 18
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine

(xi) SEQUENCE DESCRIPTION: SEQ.ID NO:3:

Glu Leu Glu Glu Leu Asn Val Pro Gly Glu Ile Val Glu Ser Leu Ser
 1 5 10 15

Ser Ser Glu Glu Ser Ile Thr Arg
 20

(2) INFORMATION FOR SEQ.ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 25
 (B) TYPE: Amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: Protein
 (ix) FEATURE:
 (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 11
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine
 (ix) FEATURE:
 (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 12
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine
 (ix) FEATURE:
 (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 13
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine
 (ix) FEATURE:
 (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 16
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine

(xi) SEQUENCE DESCRIPTION: SEQ. ID NO:4:

Asn Ala Asn Glu Glu Glu Tyr Ser Ile Gly Ser Ser Ser Glu Glu Ser
 1 5 10 15

Ala Glu Val Ala Thr Glu Glu Val Lys
 20 25

-12-

- (2) INFORMATION FOR SEQ.ID NO:5:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21
 - (B) TYPE: Amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: Protein
 - (ix) FEATURE:
 - (A) NAME/KEY: Phosphoserine
 - (B) LOCATION: 6
 - (D) OTHER INFORMATION: Post-translationally phosphorylated serine
 - (ix) FEATURE:
 - (A) NAME/KEY: Phosphoserine
 - (B) LOCATION: 8
 - (D) OTHER INFORMATION: Post-translationally phosphorylated serine
 - (ix) FEATURE:
 - (A) NAME/KEY: Phosphoserine
 - (B) LOCATION: 9
 - (D) OTHER INFORMATION: Post-translationally phosphorylated serine
 - (ix) FEATURE:
 - (A) NAME/KEY: Phosphoserine
 - (B) LOCATION: 10
 - (D) OTHER INFORMATION: Post-translationally phosphorylated serine
 - (ix) FEATURE:
 - (A) NAME/KEY: Phosphoserine
 - (B) LOCATION: 17
 - (D) OTHER INFORMATION: Post-translationally phosphorylated serine
 - (xi) SEQUENCE DESCRIPTION: SEQ. ID NO:5:
- Gln Met Glu Ala Glu Ser Ile Ser Ser Ser Glu Glu Ile Val Pro Asp
 1 5 10 15
 Ser Val Glu Gln Lys
 20

- (2) INFORMATION FOR SEQ.ID NO:6:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21
 - (B) TYPE: Amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: Protein
 - (ix) FEATURE:
 - (A) NAME/KEY: Phosphoserine
 - (B) LOCATION: 6
 - (D) OTHER INFORMATION: Post-translationally phosphorylated serine
 - (ix) FEATURE:
 - (a) Name/Key: Phosphoserine
 - (b) Location: 8
 - (d) Other information: Post-translationally phosphorylated serine
 - (ix) Feature:
 - (a) Name/Key: Phosphoserine
 - (b) Location: 9
 - (d) Other information: Post-translationally phosphorylated serine
 - (ix) Feature:
 - (a) Name/Key: Phosphoserine
 - (b) Location: 10
 - (d) Other information:

- 13 -

- Post-translationally phosphorylated serine
- (ix) Feature:
- (a) Name/Key: Phosphoserine
 - (b) Location: 17
 - (d) Other information:
Post-translationally phosphorylated serine

(xi) SEQUENCE DESCRIPTION: SEQ.ID NO:6:

Gln Met Glu Ala Glu Ser Ile Ser Ser Ser Glu Glu Ile Val Pro Asp
 1 5 10 15
 Ser Val Glu Glu Lys
 20

(2) INFORMATION FOR SEQ.ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
 - (B) TYPE: Amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Protein
- (ix) Feature:
- (a) Name/Key: Phosphoserine
 - (b) Location: 7
 - (d) Other information:
Post-translationally phosphorylated serine
- (ix) Feature:
- (a) Name/Key: Phosphoserine
 - (b) Location: 8
 - (d) Other information:
Post-translationally phosphorylated serine
- (ix) Feature:
- (a) Name/Key: Phosphoserine
 - (b) Location: 9
 - (d) Other information:
Post-translationally phosphorylated serine
- (ix) Feature:
- (a) Name/Key: Phosphoserine
 - (b) Location: 15
 - (d) Other information:
Post-translationally phosphorylated serine

(xi) SEQUENCE DESCRIPTION: SEQ.ID NO:7:

Asn Thr Met Glu His Val Ser Ser Ser Glu Glu Ser Ile Ile Ser Gln
 1 5 10 15
 Glu Thr Tyr Lys
 20

(2) INFORMATION FOR SEQ.ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 21
 - (B) TYPE: Amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Protein
- (ix) FEATURE:
- (a) Name/Key: Phosphoserine
 - (b) Location: 8
 - (d) Other information:
Post-translationally phosphorylated serine
- (ix) FEATURE:
- (a) Name/Key: Phosphoserine
 - (b) Location: 9
 - (d) Other information:

- 14 -

Post-translationally phosphorylated serine

- (ix) FEATURE:
 (a) Name/Key: Phosphoserine
 (b) Location: 10
 (d) Other information:
 Post-translationally phosphorylated serine

- (ix) FEATURE:
 (a) Name/Key: Phosphoserine
 (b) Location: 16
 (d) Other information:
 Post-translationally phosphorylated serine

(xi) SEQUENCE DESCRIPTION: SEQ.ID NO:8:

Lys Asn Thr Met Glu His Val Ser Ser Ser Glu Glu Ser Ile Ile Ser
 1 5 10 15

Gln Glu Thr Tyr Lys
 20

(2) INFORMATION FOR SEQ.ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 37
 (B) TYPE: Amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Protein

(ix) FEATURE:

- (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 4
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine

(ix) FEATURE:

- (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 6
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine

(ix) FEATURE:

- (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 22
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine

(ix) FEATURE:

- (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 24
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine

(ix) FEATURE:

- (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 25
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine

(ix) FEATURE:

- (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 26
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine

(ix) FEATURE:

- (A) NAME/KEY: Phosphoserine
 (B) LOCATION: 33
 (D) OTHER INFORMATION:
 Post-translationally phosphorylated serine

(xi) SEQUENCE DESCRIPTION: SEQ.ID NO:9:

Asp Ile Gly Ser Glu Ser Thr Glu Asp Gln Ala Met Glu Asp Ile Lys
 1 5 10 15

- 15 -

Gln Met Glu Ala Glu Ser Ile Ser Ser Ser Glu Glu Ile Val Pro Asn
20 25 30

Ser Val Glu Gln Lys
35

CLAIMS

1. A method for the preparation of selected phosphopeptides comprising the steps of completely digesting a soluble monovalent cation salt of casein in solution, introducing a di or trivalent metal ion to cause aggregation of at least the selected phosphopeptides in said digested solution, and filtering the solution through a filter having a molecular weight exclusion limit selected to retain at least said aggregated phosphopeptides while passing the bulk of the remaining phosphopeptides in a filtrate.
2. The method of claim 1, wherein the selected phosphopeptides are anticariogenic phosphopeptides and the molecular weight exclusion limit adopted during the filtering step substantially falls within the range 10,000 to 20,000.
3. The method of claim 1 or 2, wherein the soluble monovalent cation salt of casein is present in the solution in a concentration substantially falling within the range 0.1 to 50% w/w.
4. The method of any one of claims 1 to 3, wherein the digestion step is performed using a proteolytic enzyme and the ratio of proteolytic enzyme to soluble monovalent cation salt of casein in the solution falls substantially within the range 1:1000 to 1:10 (w:w)

- 17 -

selected to allow complete digestion of the casein salt.

5. The method of any preceding claim, wherein the pH and the temperature of the solution is controlled to allow complete digestion of the casein salt.

6. The method of claim 5, wherein the temperature of the solution lies substantially within the range 20°C to 60°C.

7. A method for the preparation of selected phosphopeptides having anticariogenic and other activities, comprising the steps of completely digesting a soluble monovalent cation salt of casein in solution with a proteolytic enzyme, adding a mineral acid to the solution to adjust the pH to about 4.7, removing any precipitate produced, adding CaCl₂ to a level of about 1.0% w/v to cause aggregation of at least the selected phosphopeptides in said digested solution, separating the aggregated phosphopeptides from the solution through a filter having a molecular weight exclusion limit lying substantially within the range 10,000 to 20,000 while passing the bulk of the remaining phosphopeptides and solution in a filtrate, diafiltering the separated phosphopeptides with water through a filter and concentrating and drying the retentate.

8. The method of claim 1 or 7, wherein the soluble

- 18 -

monovalent cation salt of casein is selected from sodium caseinate and potassium caseinate.

9. The method of claim 7 or 8, wherein proteolytic enzyme is selected from pancreatin, trypsin, papain, chymotrypsin or mixtures thereof.

10. A method for the preparation of anticariogenic phosphopeptides substantially as hereinbefore described with reference to the specific example.

11. Phosphopeptides when produced by the method of any preceding claim.

12. The filtrate of the method of any one of claims 1 to 10.

INTERNATIONAL SEARCH REPORT

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all)⁶

According to International Patent classification (IPC) or to both National Classification and IPC
Int. Cl.⁸ C07K 3/02, 7/10, 15/24, C12P 21/06, A23J 1/22, A23J 3/10

II. FIELDS SEARCHEDMinimum Documentation Searched⁷

Classification System

Classification Symbols

IPC⁵

C07K 3/02, 7/10, 15/24, C12P 21/06, A23J 1/22, 3/10

IPC³

C07C 103/52

Documentation Searched other than Minimum Documentation
to the extent that such Documents are included in the Fields Searched⁸

AU: IPC as above

III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹

Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate of the relevant passages ¹²	Relevant to Claim No ¹³
X	Patent Abstracts of Japan, C-260, page 49, JP, A, 59-159793 (MEIJI SEIKA K.K.) 28 February, 1983 (28.02.83).	1 to 6
X	Patent Abstracts of Japan, C261, page 66, JP, A, 59-162843 (MEIJI SEIKA K.K.) 8 March, 1983 (08.03.83).	1 to 6
X	Derwent Abstract Accession no. 76-58968X/31, Class D16, JP, A, 51-070888 (IDEMITSU KOSAN) 18 June, 1976 (18.06.76).	1
(continued)		

• Special categories of cited documents : ¹⁰

"A" Document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier document but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

"T" Later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
 "&" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search
6 July 1992 (06.07.92)

Date of Mailing of this International Search Report


16 July 1992 (16.07.92)

International Searching Authority

AUSTRALIAN PATENT OFFICE

Signature of Authorized Officer

A. BESTOW



III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category*	Citation of Document, ¹¹ with indication, where appropriate of the relevant passages ¹²	Relevant to Claim No ¹³
X	EP, A, 0090406 (MEIJI SEIKA KAISHA LTD) 5 October, 1983 (05.10.83)	1 to 6
A	See page 19, line 6 to page 22, line 1 and page 8, line 8-17.	7 to 12
X	AU, B, 51491/85 (600225) (ROUSSEL UCLAF) 12 June 1986 (12.06.86). See page 9, line 6 to page 18, line 22.	1 to 12
X	AU, B, 66783/81 (548658) (ROUSSEL UCLAF) 6 August, 1981 (06.08.81). See pages 11 to 18.	1 to 12

**ANNEX TO THE INTERNATIONAL SEARCH REPORT ON
INTERNATIONAL APPLICATION NO. PCT/AU 92/00175**

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report		Patent Family Member		
EP	90406	CA 1237937	JP 58170440	
JP	84159739	JP 61039376	US 4694821	
JP	84162843	FR 2568698	JP 61042040	US 4782440
AU	51491	AU 66783/81	CA 1178908	DK 426/81
		EP 34083	ES 498963	ES 8201408
		FI 810175	FR 2474829	JP 56123922
		JP 2257853	JP 2257854	NO 810335
		US 4358465	US 4495176	US 4740462
		US 4816398	US 5028589	ZA 8100591
AU	66783/81	CA 1178908	DK 426/81	EP 34083
		ES 498963	ES 8201408	FI 810175
		FR 2474829	JP 56123922	JP 2257853
		JP 2257854	NO 810335	US 4358465
		US 4740462	US 4816398	US 5028589
		ZA 8100591		

END OF ANNEX